

Article

Isothiourea-Catalyzed Enantioselective α -Alkylation of Esters via 1,6-Conjugate Addition to *para*-Quinone Methides

Jude N. Arokianathar ¹, Will C. Hartley ¹, Calum McLaughlin ¹, Mark D. Greenhalgh ^{1,2}, Darren Stead ³, Sean Ng ⁴, Alexandra M. Z. Slawin ¹  and Andrew D. Smith ^{1,*} 

¹ EaStCHEM, School of Chemistry, University of St Andrews, North Haugh, St Andrews KY16 9ST, UK; jude1991@hotmail.co.uk (J.N.A.); whartley@iciq.es (W.C.H.); mclaughl@uni-muenster.de (C.M.); mark.greenhalgh@warwick.ac.uk (M.D.G.); amzs@st-andrews.ac.uk (A.M.Z.S.)

² Department of Chemistry, University of Warwick, Coventry CV4 7AL, UK

³ AstraZeneca, Oncology R&D, Research & Early Development, Darwin Building, 310, Cambridge Science Park, Milton Road, Cambridge CB4 0WG, UK; Darren.Stead@astrazeneca.com

⁴ Syngenta, Jealott's Hill International Research Centre, Bracknell RG42 6EY, UK; sean.ng@syngenta.com

* Correspondence: ads10@st-andrews.ac.uk

Abstract: The isothiourea-catalyzed enantioselective 1,6-conjugate addition of *para*-nitrophenyl esters to 2,6-disubstituted *para*-quinone methides is reported. *para*-Nitrophenoxide, generated in situ from initial *N*-acylation of the isothiourea by the *para*-nitrophenyl ester, is proposed to facilitate catalyst turnover in this transformation. A range of *para*-nitrophenyl ester products can be isolated, or derivatized in situ by addition of benzylamine to give amides at up to 99% yield. Although low diastereocontrol is observed, the diastereoisomeric ester products are separable and formed with high enantiocontrol (up to 94:6 er).

Keywords: isothiourea; ammonium enolate; aryloxide; quinone methide; ester functionalization; 1,6-conjugate addition



Citation: Arokianathar, J.N.; Hartley, W.C.; McLaughlin, C.; Greenhalgh, M.D.; Stead, D.; Ng, S.; Slawin, A.M.Z.; Smith, A.D. Isothiourea-Catalyzed Enantioselective α -Alkylation of Esters via 1,6-Conjugate Addition to *para*-Quinone Methides. *Molecules* **2021**, *26*, 6333. <https://doi.org/10.3390/molecules26216333>

Academic Editors: Igor Djerdj and Alejandro Baeza Carratalá

Received: 31 August 2021

Accepted: 14 October 2021

Published: 20 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.




Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Quinone methides (QMs) are electrophilic compounds composed of a cyclohexadiene core bearing a carbonyl either *ortho* or *para* to an exocyclic alkylidene unit [1,2]. Due to their electrophilicity [3–6], QMs have been used in a variety of biological and medicinal processes [1,2,7], are present within natural products and pharmaceuticals [1,2,8–11], and have been applied as electrophiles in a variety of synthetic reactions [1,2,12–18]. While *ortho*-QMs have been used extensively in enantioselective catalysis [19], particularly as components in formal [4+2] cycloaddition reactions, the use of *para*-QMs has only recently received increased attention [19–23]. The majority of enantioselective organocatalytic methods that involve *para*-QMs have utilized Brønsted acid [24–33] or hydrogen bonding catalysts [34–38], with only a relatively small number of examples using Lewis base catalysis [39–51].

C(1)-Ammonium enolate intermediates [52–55], generated by the reaction of a tertiary amine Lewis base catalyst with a ketene, anhydride or acyl imidazole [56], have found widespread application for the synthesis of heterocyclic scaffolds in high yield and with excellent enantiocontrol. Traditionally these approaches have been limited by the requirement for the electrophilic reaction partner to contain a latent nucleophilic site to facilitate catalyst turnover. This conceptual obstacle has resulted in catalysis via C(1)-ammonium enolates being mostly applied for formal cycloaddition reactions. More recently, aryl esters have emerged as alternative C(1)-ammonium enolate precursors [55,57,58]. Significantly, following acylation of the tertiary amine catalyst by the aryl ester, a nucleophilic aryloxide is liberated, which may be exploited again in the catalytic cycle to facilitate ammonium enolate formation and catalyst turnover (Scheme 1a) [55,59]. This strategy offers a potentially

general solution to allow the expansion of electrophile scope within catalytic processes using C(1)-ammonium enolate intermediates. In 2014, we applied this concept for the isothiourea-catalyzed [2,3]-rearrangement of allylic ammonium ylides (Scheme 1b) [60–64]. More recently, this approach has been used by Snaddon (Scheme 1c) [65–72], Hartwig (Scheme 1d) [73] and Gong [74,75] for co-operative isothiourea/transition metal-catalyzed α -functionalization of pentafluorophenyl esters. In both cases, an isothiourea-derived C(1)-ammonium enolate is intercepted by an electrophilic transition metal complex to affect an allylation or benzylation reaction. We have expanded the scope of electrophiles applicable within this catalyst turnover strategy to include iminium ions generated under either photoredox conditions or Brønsted acid catalysis, as well as bis-sulfone Michael acceptors, and pyridinium salts (Scheme 1e) [76–79]. Recently, Waser also reported an elegant example of this turnover strategy for the enantioselective α -chlorination of pentafluorophenyl esters [80], while Zheng and co-workers reported a related approach using diphenyl methanol as an external turnover reagent for the fluorination of carboxylic acids [81,82]. A significant challenge within this area is the identification of electrophilic reaction partners that react with the catalytically-generated C(1)-ammonium enolate, but are compatible with the nucleophilic tertiary amine catalyst and aryloxide, which is essential for catalyst turnover. Building upon this conceptual platform, it was envisaged that *para*-QMs may be suitable electrophiles to apply in formal 1,6-conjugate additions.



Scheme 1. Isothiourea-catalyzed enantioselective processes using aryloxide-facilitated catalyst turnover.


2. Results

2.1. Reaction Optimization

Initial studies focused on the isothiourea-catalyzed 1,6-conjugate addition of *para*-nitrophenyl (PNP) ester **1** to 2,6-di-*tert*-butyl *para*-QM **5** (Table 1). Benzylamine was added at the end of the reaction to convert the PNP ester product to the corresponding amide. Based on our previous experience, the amide was expected to be more stable to chromatographic purification. Using tetramisole HCl **6** (20 mol%) as the catalyst, *i*-Pr₂NEt as the base and MeCN as the solvent gave a 55:45 ratio of diastereoisomeric amide products **7** and **8** in high yield (82%) and with excellent enantioselectivity (**7**: 97:3 er; **8**: 94:6 er) (Entry 1). Chromatographic separation of the diastereoisomers was not possible, however, the enantioenrichment of both **7** and **8** could be reliably determined by chiral stationary phase (CSP)-HPLC analysis of the mixture. A control reaction in the absence of the catalyst showed no conversion (Entry 2). The use of six alternative solvents was investigated (PhMe, CH₂Cl₂, CHCl₃, THF, 1,4-dioxane and DMF), (see the Supporting Information for details) however, only the use of DMF provided any conversion to the product, indicating that solvent polarity may be significant for the success of this transformation. A control reaction in DMF in the absence of the catalyst, however, also led to comparable conversion to the product, consistent with the operation of a competitive Brønsted base-promoted reaction (see the Supporting Information for details). Taking MeCN as the optimal solvent, the use of eight different organic and inorganic bases was investigated (see the Supporting Information for details). Of those tested, Et₃N provided an improved yield of 98%, whilst maintaining comparable diastereo- and enantioselectivity (Entry 3). The use of alternative aryl esters was next probed, with pentafluorophenyl ester **2** and bis(trifluoromethyl)phenyl ester **3** giving amide products **7** and **8** in high yield, but with lower enantioselectivity than when using PNP ester **1** (Entries 4 and 5). The use of 2,4,6-trichlorophenyl ester **4** resulted in only 31% yield (Entry 6), which is consistent with previous studies in this field [65,72,76,79], and most likely reflects the increased steric hindrance of the aryloxide attenuating its nucleophilicity. Finally, using PNP ester **1**, the catalyst loading could be reduced to 5 mol% with only a small drop in stereoselectivity (Entry 7), while heating the reaction to 40 °C provided a slight improvement in yield (Entry 8). The reaction could also be performed in the absence of a base (Entry 9), however, slightly lower yield was obtained and therefore, during investigation of the substrate scope, Et₃N was routinely used as an auxiliary base.

2.2. Reaction Scope and Limitations


Due to the low diastereoselectivity observed using 2,6-di-*tert*-butyl *para*-QM **5**, the alternative use of 2,6-disubstituted *para*-QMs were investigated (Scheme 2). 2,6-Dimethyl, dibromo and diphenyl *para*-QMs **9–11** bearing a phenyl substituent at the exocyclic olefin were applied under optimized conditions. In all cases significantly lower conversion was observed ($\leq 43\%$), and the amide products **14–16** were difficult to isolate. Based on analysis of the crude reaction mixture by ¹H NMR spectroscopy, the 2,6-dimethyl and dibromo-substituted analogues **14** and **15** were obtained with marginally improved diastereoselectivity (~70:30 dr), indicating that alternative substituents in these positions could prove beneficial if the products were isolable. Next, variation of the exocyclic substituent was probed. Incorporation of a methyl group at this position provided no improvement in dr, but the amide product **17** was isolated in a 50% yield and with moderate enantioenrichment for both diastereoisomers. Finally, incorporation of a 2-naphthyl substituent at this position was well tolerated, with amide **18** obtained in an 83% yield, 70:30 dr and excellent enantiocontrol (94:6 er) for the major diastereoisomer.

Table 1. Reaction optimization.


Entry	6 (mol%)	Aryl Ester	Base	Yield (%)	dr (7:8)	er (7)	er (8)
1	20	1	<i>i</i> -Pr ₂ NEt	82	55:45	97:3	94:6
2	0	1	<i>i</i> -Pr ₂ NEt	0	-	-	-
3	20	1	Et ₃ N	98	60:40	97:3	94:6
4	20	2	Et ₃ N	99	65:35	81:19	94:6
5	20	3	Et ₃ N	88	60:40	94:6	91:9
6	20	4	Et ₃ N	31	55:45	84:16	71:29
7	5	1	Et ₃ N	95	55:45	93:7	88:12
8 [a]	5	1	Et ₃ N	99	60:40	92:8	85:15
9 [a]	5	1	none	90	60:40	93:7	89:11

All reactions were carried out on a 0.25 mmol scale; isolated yields are a mixture of diastereoisomers **7** and **8**; dr was determined by ¹H NMR spectroscopic analysis of the crude reaction mixture; er was determined by CSP-HPLC analysis: **7** (2*S*,1'*R*:2*R*,1'*S*) and **8** (2*S*,1'*S*:2*R*,1'*R*). [a] Reaction performed at 40 °C.

Although structural variation of the *para*-QM provided marginal improvements in dr, the use of 2,6-di-*tert*-butyl *para*-QM **5** was considered most convenient for further investigations due to its stability and ease of synthesis, and the higher yields of product obtained from catalysis. To investigate if the dr obtained in these reactions was a manifestation of a kinetic or thermodynamic preference, isolation of diastereoisomeric PNP esters **19** and **20** and resubjection to catalysis conditions was attempted. Although the diastereoisomeric amides **7** and **8** had proved difficult to separate, the corresponding PNP esters **19** and **20** were chromatographically separable and displayed high stability. Epimerization studies were conducted using each diastereoisomer through sequential treatment with *i*-Pr₂NEt, (S)-TM HCl **6**, *para*-nitrophenoxide and benzylamine (Scheme 3). These experiments were followed by in situ ¹H NMR spectroscopic analysis and revealed no epimerization in either case. This indicates the dr obtained in the catalytic reaction most likely reflects the inherent diastereoselectivity of the transformation. Following separation of the diastereoisomers, the absolute configuration of the major diastereoisomer could also be confirmed as (2*S*,1'*R*) by single crystal X-ray crystallographic analysis [83,84]. Based on literature precedent, the (S)-configuration at C(2) was expected to be generated under catalyst-control, and therefore the absolute configuration of the minor diastereoisomer was predicted to be (2*S*,1'*S*).




Scheme 2. Scope: Variation of *para*-quinone methide—0.25 mmol scale; only the structure of the major diastereoisomer is shown; isolated yield is a mixture of diastereoisomers; dr determined by ^1H NMR spectroscopic analysis of the crude reaction product mixture; er determined by CSP-HPLC analysis. ^[a] Product not isolable: conversion based on ^1H NMR analysis of the crude reaction product mixture; ers could not be determined.



Scheme 3. Control studies and confirmation of absolute configuration of products. The majority of hydrogen atoms are omitted for clarity within X-ray crystal structure representation of **19**.


Having established the stability of the PNP ester products and demonstrated the potential to separate the diastereoisomers by column chromatography, the scope of the catalytic transformation was investigated through variation of the PNP ester substrate. In each case, the PNP ester product diastereoisomers were at least partially separable, enabling unambiguous characterization. To test the applicability of the procedure, *p*-tolyl-substituted PNP ester product **19** was prepared on a larger scale (1.25 mmol) (Scheme 4). A combined 85% yield of both diastereoisomers was obtained, with comparable stereoselectivity to that observed when the reaction was conducted on an analytical scale (Table 1, Entry 7). The generality of the procedure was further probed using five electronically- and sterically-differentiated PNP esters. Introduction of an electron-donating 4-methoxy substituent was well tolerated, with PNP ester **26** obtained in quantitative yield, 60:40 dr and with high enantioselectivity for both diastereoisomers. Under the optimized conditions, the introduction of an electron-withdrawing 4-trifluoromethyl group resulted in low enantioselectivity (**27**: 63:37 er), which was attributed to a competitive Brønsted base-promoted background reaction. Consistent with this hypothesis, repeating the reaction in the absence of Et₃N, and using the free base of the isothioureia catalyst **6**, provided PNP ester product **27** in 94% yield and significantly improved enantioselectivity (88:12 er_{major}; 79:21 er_{minor}). A similar effect was observed when using a 2-naphthyl-substituted PNP ester, with optimal enantioselectivity obtained in the absence of an auxiliary base (**28**: 91:9 er_{major}; 85:15 er_{minor}). Introduction of a sterically-imposing 1-naphthyl or a heteroaromatic thienyl substituent was also tolerated, with **29** and **30** obtained in excellent yield and with high enantioselectivity.



Scheme 4. Scope: Variation of *para*-nitrophenyl ester—0.25 mmol scale; only the structure of the major diastereoisomer is shown; isolated yields are given for the combination of diastereoisomers; dr determined by ¹H NMR spectroscopic analysis of the crude reaction product mixture; er determined by CSP-HPLC analysis. ^[a] 1.25 mmol scale. ^[b] Conducted at r.t., in the absence of Et₃N, and using free base of **6**.

2.3. Proposed Mechanism

The mechanism of this transformation is proposed to begin with *N*-acylation of the free base isothiourea catalyst **6** by PNP ester **31** to generate the corresponding acyl ammonium *para*-nitrophenoxide ion pair **32** (Scheme 5). Subsequent deprotonation leads to (Z)-ammonium enolate **33**. Based on previous mechanistic studies [78,85], and the catalytic activity observed in the absence of an auxiliary base (Table 1, Entry 9), deprotonation is likely to be affected by the *para*-nitrophenoxide counterion. 1,6-Conjugate addition of ammonium enolate **33** to *para*-QM electrophile **34**, followed by protonation, gives acyl ammonium intermediate **35**. Finally, regeneration of catalyst **6**, and concurrent release of product **37**, is proposed to be facilitated by *para*-nitrophenoxide [59–79,86,87]. Although not essential for reactivity, the addition of Et₃N as an auxiliary base may be beneficial as a proton shuttle, and to maintain the isothiourea catalyst in its non-protonated form **6** [77,86]. The enantioselectivity of the transformation indicates the C–C bond forming event takes place on the *Si*-face of the ammonium enolate. This selectivity can be rationalized through preferential formation of the (Z)-ammonium enolate [76–79,85], which is conformationally-restricted by an intramolecular 1,5-O⋯S interaction [61,88–104] and results in the phenyl stereodirecting group of the catalyst blocking the enolate *Re*-face. The observed poor diastereoselectivity can be tentatively rationalized by a simple stereochemical model that assumes a favored, open pre-transition state assembly where steric interactions are minimized about the forming C–C bond. Minimal differentiation between the aryl- and quinone substituents of the *para*-QM quinone leads to the two transition state assemblies **38** and **39** that give the major and minor diastereoisomers, respectively.



Scheme 5. Proposed mechanism (only the pathway for the formation of the major diastereoisomer is shown) and transition state assemblies. ArOH corresponds to either PNPOH or the reaction product.

3. Materials and Methods

3.1. General Procedure for the Enantioselective 1,6-Addition

In a flame-dried vial, the requisite *para*-quinone methide (1.0 equiv.), aryl ester (1.5 equiv.), (S)-TM HCl (5 mol%), Et₃N (1.0 equiv.) and anhydrous MeCN (0.6 M) was added and stirred at r.t. for 24 h. The reaction was then quenched with benzylamine (5.0 equiv.) and stirred at r.t. for a further 12 h before being concentrated in vacuo. The residue was diluted with EtOAc (20 mL) and washed successively with 10% citric acid (20 mL × 1), aqueous NaOH (20 mL × 3) and brine (20 mL × 1). The organic layer was extracted, dried over MgSO₄ and the filtrate was concentrated in vacuo. The crude material was purified by flash silica column chromatography to give the desired product.

3.2. Representative Synthesis and Characterization of Compounds 7 and 8 (Entry 8)

Following the general procedure above, 4-benzylidene-2,6-di-*tert*-butylcyclohexa-2,5-dien-1-one **5** (74 mg, 0.25 mmol), 4'-nitrophenyl 2-(*p*-tolyl)acetate **1** (102 mg, 0.375 mmol), (S)-TM HCl **6** (3 mg, 5 mol%) and Et₃N (35 µL, 0.25 mmol) were dissolved in anhydrous MeCN (0.42 mL). The reaction mixture was stirred at 40 °C for 24 h before being quenched with benzylamine (137 µL, 1.25 mmol) at r.t. to give a crude mixture containing the title compound in 60:40 dr. The mixture was purified by flash silica column chromatography (petroleum ether/EtOAc, 85:15) to afford diastereoisomers **7** and **8** (60:40 dr) (132 mg, 99%) as a pale yellow solid.

mp 126–128 °C; $[\alpha]_D^{20}$ +10.0 (*c* 1.0, CHCl₃); IR ν_{\max} (film)/cm^{−1} 3638 (O–H) 3304 (N–H), 2955 (C–H), 1645 (C=O); HRMS (ESI⁺) C₃₇H₄₃NO₂ ([M + H]⁺), found 534.3359, requires 534.3367 (−1.4 ppm).

Data for major diastereoisomer (7): Chiral HPLC analysis, Chiralpak AD-H (10% *i*-PrOH/hexane, flow rate 1.5 mLmin^{−1}, 211 nm, 40 °C), *t*_R 8.5 min and 29.3 min, 92:8 er; ¹H NMR (500 MHz, CDCl₃) δ_H : 1.42 (18H, s, (C(3'''))C(CH₃)₃, C(5'''))C(CH₃)₃), 2.24 (3H, s, C(4'')CH₃), 3.90–4.05 (2H, m, C(2)H, CH_APh), 4.44 (1H, dd, *J* 15.0, 6.8, CH_BPh), 4.82 (1H, d, *J* 11.7, C(1')H), 5.14 (1H, s, OH), 5.55 (1H, t, *J* 5.6, NH), 6.71–6.76 (2H, m, *Ar*), 6.96–7.05 (2H, m, *Ar*), 7.09–7.15 (2H, m, *Ar*), 7.15–7.22 (4H, m, *Ar*), 7.23–7.30 (4H, m, C(4'')H, C(2''')H, C(6''')H, *Ar*), 7.34 (1H, t, *J* 7.6, *Ar*), 7.43–7.51 (1H, m, *Ar*); ¹³C{¹H} NMR (126 MHz, CDCl₃) δ_C : 21.0 (C(4'')CH₃), 30.4 (C(3'''))C(CH₃)₃, C(5'''))C(CH₃)₃, 34.4 (C(3'''))C(CH₃)₃, C(5'''))C(CH₃)₃, 43.6 (CH₂Ph), 54.1 (C(1')), 59.5 (C(2)), 124.7 (C(2'''), C(6''')), 125.8 (C(4'')), 128.2 (*Ar*), 127.2 (*Ar*), 127.3 (*Ar*), 128.0 (*Ar*), 128.4 (*Ar*), 128.5 (*Ar*), 128.6 (*Ar*), 129.0 (*Ar*), 133.8 (C(1''')), 135.2 (C(1'')), 135.6 (C(3'''), C(5''')), 136.4 (C(4'')), 137.9 (*i*-Ph), 142.4 (C(1''')), 152.4 (C(4''')), 172.1 (C(1)).

Selected data for minor diastereoisomer (8): Chiral HPLC analysis, Chiralpak AD-H (10% *i*-PrOH/hexane, flow rate 1.5 mLmin^{−1}, 211 nm, 40 °C), *t*_R 3.8 min and 18.6 min, 85:15 er; ¹H NMR (500 MHz, CDCl₃) δ_H : 1.27 (18H, s, (C(3'''))C(CH₃)₃, C(5'''))C(CH₃)₃), 2.27 (3H, s, C(4'')CH₃), 3.90–4.05 (2H, m, C(2)H, CH_APh), 4.50 (1H, dd, *J* 15.0, 6.8, CH_BPh), 4.69 (1H, d, *J* 11.7, C(1')H), 4.89 (1H, s, OH), 5.69 (1H, t, *J* 5.6, NH), 6.81–6.84 (2H, m, *Ar*); ¹³C{¹H} NMR (126 MHz, CDCl₃) δ_C : 21.0 (C(4'')CH₃), 30.2 (C(3'''))C(CH₃)₃, C(5'''))C(CH₃)₃, 34.2 (C(3'''))C(CH₃)₃, C(5'''))C(CH₃)₃, 43.4 (CH₂Ph), 54.7 (C(1')), 59.5 (C(2)), 125.3 (C(2'''), C(6''')), 126.3 (C(4'')), 127.5 (*Ar*), 128.2 (*Ar*), 128.4 (*Ar*), 128.8 (*Ar*), 132.1 (C(1''')), 134.9 (C(3'''), C(5''')), 135.5 (C(1'')), 136.4 (C(4'')), 138.2 (*i*-Ph), 143.6 (C(1''')), 151.7 (C(4''')), 172.0 (C(1)).

4. Conclusions

An isothioureia-catalyzed enantioselective 1,6-conjugate addition of *para*-nitrophenyl (PNP) esters to *para*-quinone methides (QMs) has been developed. Variation of the arylacetic ester and *para*-QM substrates has provided a range of functionalized products in generally excellent yields and high enantiocontrol (up to 94:6 er). An inherent limitation of the method is that the products were routinely obtained in ~60:40 dr. This diastereoselectivity was shown to arise from kinetic control, but was relatively insensitive to changes in reaction conditions and structural variation of the substrates. Although the dr could

not be improved, the diastereoisomeric PNP ester products could be separated by column chromatography. The success of this catalytic methodology is proposed to rely upon the *para*-nitrophenoxide, expelled during *N*-acylation of the catalyst, to facilitate catalyst turnover and release the product. Current work in our laboratory is focused on further applications of using in situ-generated aryloxides to promote catalyst turnover in Lewis base catalysis.

Supplementary Materials: Full experimental procedures, characterization data, NMR spectra and HPLC chromatograms for all new compounds, as well as crystallographic data for product **19** (CCDC 1992504) are available online.

Author Contributions: Conceptualization, A.D.S. and J.N.A.; investigation, J.N.A.; W.C.H.; C.M.; X-ray crystallographic analysis, A.M.Z.S.; writing—original draft preparation, M.D.G.; writing—review and editing, A.D.S. and all authors; supervision, D.S., S.N. and A.D.S.; funding acquisition, A.D.S. All authors have read and agreed to the published version of the manuscript.

Funding: We thank the ERC under the European Union's Seventh Framework Programme (FP7/2007-2013)/E.R.C. grant agreement n° 279850, AstraZeneca and EPSRC (EP/M506631/1 (J.N.A.)), Syngenta and the EPSRC Centre for Doctoral Training in Critical Resource Catalysis (CRITICAT, EP/L016419/1 (W.C.H.)), and EPSRC (EP/M508214/1 (C.M.)) for funding. A.D.S. thanks the Royal Society for a Wolfson Research Merit Award. We thank the EPSRC UK National Mass Spectrometry Facility at Swansea University.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The research data underpinning this publication can be found at DOI: 10.17630/f6cf6c80-483d-4f16-bd79-80e1537513b2.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are available from the authors on request.

References and Notes

1. Rokita, S.E. *Quinone Methides*; Wiley: Hoboken, NJ, USA, 2009.
2. Toteva, M.M.; Richard, J.P. The generation and reactions of quinone methides. *Adv. Phys. Org. Chem.* **2011**, *45*, 39–91. [PubMed]
3. Turner, A.B. Quinone methides. *Q. Rev. Chem. Soc.* **1964**, *18*, 347–360. [CrossRef]
4. Wagner, H.-U.R. Gompper. In *The Chemistry of Quinonoid Compounds*; Wiley: London, UK, 1974; pp. 1145–1178.
5. Richter, D.; Hampel, N.; Singer, T.; Ofial, A.R.; Mayr, H. Synthesis and Characterization of Novel Quinone Methides: Reference Electrophiles for the Construction of Nucleophilicity Scales. *Eur. J. Org. Chem.* **2009**, 3203–3211. [CrossRef]
6. Singh, M.S. *Reactive Intermediates in Organic Chemistry: Structure and Mechanism*; Wiley-VCH: Weinheim, Germany, 2014.
7. Freccero, M. Quinone methides as alkylating and cross-linking agents. *Mini Rev. Org. Chem.* **2004**, *1*, 403–415. [CrossRef]
8. Peters, M.G. Chemical modifications of biopolymers by quinones and quinone methides. *Angew. Chem. Int. Ed.* **1989**, *28*, 555–570, *Angew. Chem.* **1989**, *101*, 572–587. [CrossRef]
9. Martin, H.J.; Magauer, T.; Mulzer, J. In Pursuit of a Competitive Target: Total Synthesis of the Antibiotic Kendomycin. *Angew. Chem. Int. Ed.* **2010**, *49*, 5614–5626, *Angew. Chem.* **2010**, *122*, 5746–5758. [CrossRef]
10. Jansen, R.; Gerth, K.; Steinmetz, H.; Reinecke, S.; Kessler, W.; Kirschning, A.; Müller, R. Elansolid A3, a Unique p-Quinone Methide Antibiotic from *Chitinophaga sancti*. *Chem. Eur. J.* **2011**, *17*, 7739–7744. [CrossRef]
11. Dehn, R.; Katsuyama, Y.; Weber, A.; Gerth, K.; Jansen, R.; Steinmetz, H.; Höfle, G.; Müller, R.; Kirschning, A. Molecular Basis of Elansolid Biosynthesis: Evidence for an Unprecedented Quinone Methide Initiated Intramolecular Diels–Alder Cycloaddition/Macrolactonization. *Angew. Chem. Int. Ed.* **2011**, *50*, 3882–3887, *Angew. Chem.* **2011**, *123*, 3968–3973. [CrossRef]
12. Gai, K.; Fang, X.; Li, X.; Xu, J.; Wu, X.; Lin, A.; Yao, H. Synthesis of spiro[2.5]octa-4,7-dien-6-one with consecutive quaternary centers via 1,6-conjugate addition induced dearomatization of para-quinone methides. *Chem. Commun.* **2015**, *51*, 15831–15834. [CrossRef]
13. López, A.; Parra, A.; Jarava-Barrera, C.; Tortosa, M. Copper-catalyzed silylation of p-quinone methides: New entry to dibenzylic silanes. *Chem. Commun.* **2015**, *51*, 17684–17687. [CrossRef]
14. Ramanjaneyulu, B.T.; Mahesh, S.; Anand, R.V. Bis(amino)cyclopropenylidene-Catalyzed 1,6-Conjugate Addition of Aromatic Aldehydes to para-Quinone Methides: Expedient Access to α,α' -Diarylated Ketones. *Org. Lett.* **2015**, *17*, 3952–3955. [CrossRef]
15. Yuan, Z.; Fang, X.; Li, X.; Wu, J.; Yao, H.; Lin, A. 1,6-Conjugated Addition-Mediated [2+1] Annulation: Approach to Spiro[2.5]octa-4,7-dien-6-one. *J. Org. Chem.* **2015**, *80*, 11123–11130. [CrossRef] [PubMed]

16. Reddy, V.; Anand, R.V. Expedient Access to Unsymmetrical Diarylindolylmethanes through Palladium-Catalyzed Domino Electrophilic Cyclization–Extended Conjugate Addition Approach. *Org. Lett.* **2015**, *17*, 3390–3393. [\[CrossRef\]](#)
17. Roiser, L.; Zielke, K.; Waser, M. Enantioselective Spirocyclopropanation of para-Quinone Methides using Ammonium Ylides. *Org. Lett.* **2017**, *19*, 2338–2341. [\[CrossRef\]](#)
18. Roiser, L.; Zielke, K.; Waser, M. Formal (4+1) Cyclization of Ammonium Ylides with Vinylogous para-Quinone Methides. *Synthesis* **2018**, *50*, 4047–4054.
19. Caruana, L.; Fochi, M.; Bernardi, L. The Emergence of Quinone Methides in Asymmetric Organocatalysis. *Molecules* **2015**, *20*, 11733–11764. [\[CrossRef\]](#)
20. Parra, A.; Tortosa, M. para-Quinone Methide: A New Player in Asymmetric Catalysis. *ChemCatChem* **2015**, *7*, 1524–1526. [\[CrossRef\]](#)
21. Chauhan, P.; Kaya, U.; Enders, D. Advances in Organocatalytic 1,6-Addition Reactions: Enantioselective Construction of Remote Stereogenic Centers. *Adv. Synth. Catal.* **2017**, *359*, 888–912. [\[CrossRef\]](#)
22. Li, W.; Xu, X.; Zhang, P.; Li, P. Recent Advances in the Catalytic Enantioselective Reactions of para-Quinone Methides. *Chem. Asian J.* **2018**, *13*, 2350–2359. [\[CrossRef\]](#)
23. Wang, J.-Y.; Hao, W.-J.; Tu, S.-J.; Jiang, B. Recent developments in 1,6-addition reactions of para-quinone methides (p-QMs). *Org. Chem. Front.* **2020**, *7*, 1743–1778. [\[CrossRef\]](#)
24. Wang, Z.; Wong, Y.F.; Sun, J. Catalytic Asymmetric 1,6-Conjugate Addition of para-Quinone Methides: Formation of All-Carbon Quaternary Stereocenters. *Angew. Chem. Int. Ed.* **2015**, *54*, 13711–13714. *Angew. Chem.* **2015**, *127*, 13915–13918. [\[CrossRef\]](#)
25. Dong, N.; Zhang, Z.-P.; Xue, X.-S.; Li, X.; Cheng, J.-P. Phosphoric Acid Catalyzed Asymmetric 1,6-Conjugate Addition of Thioacetic Acid to para-Quinone Methides. *Angew. Chem. Int. Ed.* **2016**, *55*, 1460–1464. *Angew. Chem.* **2016**, *128*, 1482–1486. [\[CrossRef\]](#)
26. Wong, Y.F.; Wang, Z.; Sun, J. Chiral phosphoric acid catalyzed asymmetric addition of naphthols to para-quinone methides. *Org. Biomol. Chem.* **2016**, *14*, 5751–5754. [\[CrossRef\]](#)
27. Chen, M.; Sun, J. How understanding the role of an additive can lead to an improved synthetic protocol without an additive: Organocatalytic synthesis of chiral diarylmethyl alkynes. *Angew. Chem.* **2017**, *129*, 12128–12132. *Angew. Chem. Int. Ed.* **2017**, *56*, 11966–11970. [\[CrossRef\]](#)
28. Yan, J.; Chen, M.; Sung, H.H.-Y.; Williams, I.D.; Sun, J. An Organocatalytic Asymmetric Synthesis of Chiral β,β -Diaryl- α -amino Acids via Addition of Azlactones to In Situ Generated para-Quinone Methides. *Chem. Asian J.* **2018**, *13*, 2440–2444. [\[CrossRef\]](#)
29. Rahman, A.; Zhou, Q.; Lin, X. Asymmetric organocatalytic synthesis of chiral 3,3-disubstituted oxindoles via a 1, 6-conjugate addition reaction. *Org. Biomol. Chem.* **2018**, *16*, 5301–5309. [\[CrossRef\]](#)
30. Li, W.; Xu, X.; Liu, Y.; Gao, H.; Cheng, Y.; Li, P. Enantioselective Organocatalytic 1,6-Addition of Azlactones to para-Quinone Methides: An Access to α,α -Disubstituted and β,β -Diaryl- α -amino acid Esters. *Org. Lett.* **2018**, *20*, 1142–1145. [\[CrossRef\]](#) [\[PubMed\]](#)
31. Wang, J.-R.; Jiang, X.-L.; Hang, Q.-Q.; Zhang, S.; Mei, G.-J.; Shi, F. Catalytic Asymmetric Conjugate Addition of Indoles to para-Quinone Methide Derivatives. *J. Org. Chem.* **2019**, *84*, 7829–7839. [\[CrossRef\]](#)
32. Wang, Z.; Zhu, Y.; Pan, X.; Wang, G.; Liu, L. Synthesis of Chiral Triarylmethanes Bearing All-Carbon Quaternary Stereocenters: Catalytic Asymmetric Oxidative Cross-Coupling of 2,2-Diarylacetonitriles and (Hetero)arenes. *Angew. Chem.* **2020**, *132*, 3077–3081, *Angew. Chem.* **2020**, *59*, 3053–3057. [\[CrossRef\]](#)
33. Niu, J.-P.; Nie, J.; Li, S.; Ma, J.-A. Organocatalytic asymmetric synthesis of β,β -diaryl ketones via one-pot tandem dehydration/1,6-addition/decarboxylation transformation of β -keto acids and 4-hydroxybenzyl alcohols. *Chem. Commun.* **2020**, *56*, 8687–8690. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Zhao, K.; Zhi, Y.; Wang, A.; Enders, D. Asymmetric Organocatalytic Synthesis of 3-Diarylmethine-Substituted Oxindoles Bearing a Quaternary Stereocenter via 1,6-Conjugate Addition to para-Quinone Methides. *ACS Catal.* **2016**, *6*, 657–660. [\[CrossRef\]](#)
35. Li, X.; Xu, X.; Wei, W.; Lin, A.; Yao, H. Organocatalyzed Asymmetric 1,6-Conjugate Addition of para-Quinone Methides with Dicyanoolefins. *Org. Lett.* **2016**, *18*, 428–431. [\[CrossRef\]](#)
36. Deng, Y.-H.; Zhang, X.-Z.; Yu, K.-Y.; Yan, X.; Du, J.-Y.; Huang, H.; Fan, C.-A. Bifunctional tertiary amine-squaramide catalyzed asymmetric catalytic 1,6-conjugate addition/aromatization of para-quinone methides with oxindoles. *Chem. Commun.* **2016**, *52*, 4183–4186. [\[CrossRef\]](#) [\[PubMed\]](#)
37. Torà, R.; Vila, C.; Sanz-Marco, A.; Muñoz, M.C.; Pedro, J.R.; Blay, G. Organocatalytic Enantioselective 1,6-aza-Michael Addition of Isoxazolin-5-ones to p-Quinone Methides. *Eur. J. Org. Chem.* **2020**, *2020*, 627–630. [\[CrossRef\]](#)
38. Wang, L.; Yang, F.; Xua, X.; Jiang, J. Organocatalytic 1,6-hydrophosphination of para-quinone methides: Enantioselective access to chiral 3-phosphoxindoles bearing phosphorus-substituted quaternary carbon stereocenters. *Org. Chem. Front.* **2021**, *8*, 2002–2008. [\[CrossRef\]](#)
39. Caruana, L.; Kniep, F.; Johansen, T.K.; Poulsen, H.P.; Jørgensen, K.A. A New Organocatalytic Concept for Asymmetric α -Alkylation of Aldehydes. *J. Am. Chem. Soc.* **2014**, *136*, 15929–15932. [\[CrossRef\]](#)
40. Li, S.; Liu, Y.; Huang, B.; Zhou, T.; Tao, H.; Xiao, Y.; Liu, L.; Zhang, J. Phosphine-Catalyzed Asymmetric Intermolecular Cross-Vinylogous Rauhut–Currier Reactions of Vinyl Ketones with para-Quinone Methides. *ACS Catal.* **2017**, *7*, 2805–2809. [\[CrossRef\]](#)
41. Kang, T.-C.; Wu, L.-P.; Yu, Q.-W.; Wu, X.-Y. Enantioselective Rauhut–Currier-Type 1,6-Conjugate Addition of Methyl Vinyl Ketone to para-Quinone Methides. *Chem. Eur. J.* **2017**, *23*, 6509–6513. [\[CrossRef\]](#)

42. Wang, D.; Song, Z.-F.; Wang, W.-J.; Xu, T. Highly Regio- and Enantioselective Dienylation of p-Quinone Methides Enabled by an Organocatalyzed Isomerization/Addition Cascade of Allenates. *Org. Lett.* **2019**, *21*, 3963–3967. [\[CrossRef\]](#)
43. Li, W.; Yuan, H.; Liu, Z.; Zhang, Z.; Cheng, Y.; Li, P. NHC-Catalyzed Enantioselective [4+3] Cycloaddition of Ortho-Hydroxyphenyl Substituted Para-Quinone Methides with Isatin-Derived Enals. *Adv. Synth. Catal.* **2018**, *360*, 2460–2464. [\[CrossRef\]](#)
44. Liu, Q.; Chen, X.-Y.; Rissanen, K.; Ender, D. Asymmetric synthesis of spiro-oxindole- ϵ -lactones through N-heterocyclic carbene catalysis. *Org. Lett.* **2018**, *20*, 3622–3626. [\[CrossRef\]](#)
45. Zhao, M.-X.; Xiang, J.; Zhao, Z.-Q.; Zhao, X.-L.; Shi, M. Asymmetric synthesis of dihydrocoumarins via catalytic sequential 1,6-addition/transesterification of α -isocyanoacetates with para-quinone methides. *Org. Biomol. Chem.* **2020**, *18*, 1637–1646. [\[CrossRef\]](#) [\[PubMed\]](#)
46. Roy, S.; Pradhan, S.; Kumar, K.; Chatterjee, I. Asymmetric organocatalytic double 1,6-addition: Rapid access to chiral chromans with molecular complexity. *Org. Chem. Front.* **2020**, *7*, 1388–1394. [\[CrossRef\]](#)
47. Chu, W.D.; Zhang, L.F.; Bao, X.; Zhao, X.H.; Zeng, C.; Du, J.Y.; Zhang, G.B.; Wang, F.X.; Ma, X.Y.; Fan, C.A. Asymmetric Catalytic 1,6-Conjugate Addition/Aromatization of para-Quinone Methides: Enantioselective Introduction of Functionalized Diarylmethine Stereogenic Centers. *Angew. Chem. Int. Ed.* **2013**, *52*, 9229–9233. *Angew. Chem.* **2013**, *125*, 9399–9403. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Zhang, X.-Z.; Deng, Y.-H.; Yan, X.; Yu, K.-Y.; Wang, F.-X.; Ma, X.-Y.; Fan, C.-A. Diastereoselective and Enantioselective Synthesis of Unsymmetric β , β -Diaryl- α -Amino Acid Esters via Organocatalytic 1,6-Conjugate Addition of para-Quinone Methides. *J. Org. Chem.* **2016**, *81*, 5655–5662. [\[CrossRef\]](#)
49. Ge, L.; Lu, X.; Cheng, C.; Chen, J.; Cao, W.; Wu, X.; Zhao, G. Amide-Phosphonium Salt as Bifunctional Phase Transfer Catalyst for Asymmetric 1,6-Addition of Malonate Esters to para-Quinone Methides. *J. Org. Chem.* **2016**, *81*, 9315–9325. [\[CrossRef\]](#)
50. Santra, S.; Porey, A.; Jana, B.; Guin, J. N-Heterocyclic carbenes as chiral Brønsted base catalysts: A highly diastereo- and enantioselective 1,6-addition reaction. *Chem. Sci.* **2018**, *9*, 6446–6450. [\[CrossRef\]](#)
51. Eitzinger, A.; Winter, M.; Schörgenhumer, J.; Waser, M. Quaternary β 2, 2-amino acid derivatives by asymmetric addition of isoxazolidin-5-ones to para-quinone methides. *Chem. Commun.* **2020**, *56*, 579–582. [\[CrossRef\]](#)
52. Van, K.N.; Morrill, L.C.; Smith, A.D.; Romo, D. *Lewis Base Catalysis in Organic Synthesis*; Vedejs, E., Denmark, S.E., Eds.; Wiley-VCH: Weinheim, Germany, 2016; Chapter 13; pp. 527–653.
53. Morrill, L.C.; Smith, A.D. Organocatalytic Lewis base functionalisation of carboxylic acids, esters and anhydrides via C1-ammonium or azolium enolates. *Chem. Soc. Rev.* **2014**, *43*, 6214–6226. [\[CrossRef\]](#)
54. Gaunt, M.J.; Johansson, C.C.C. Recent developments in the use of catalytic asymmetric ammonium enolates in chemical synthesis. *Chem. Rev.* **2007**, *107*, 5596–5605. [\[CrossRef\]](#)
55. McLaughlin, C.; Smith, A.D. Generation and Reactivity of C (1)-Ammonium Enolates by Using Isothiourea Catalysis. *Chem. Eur. J.* **2021**, *27*, 1533–1555. [\[CrossRef\]](#) [\[PubMed\]](#)
56. Young, C.M.; Stark, D.G.; West, T.H.; Taylor, J.E.; Smith, A.D. Exploiting the Imidazolium Effect in Base-free Ammonium Enolate Generation: Synthetic and Mechanistic Studies. *Angew. Chem.* **2016**, *128*, 14606–14611. *Angew. Chem.* **2016**, *55*, 14394–14399. [\[CrossRef\]](#) [\[PubMed\]](#)
57. Hao, L.; Chen, X.; Chen, S.; Jiang, K.; Torres, J.; Chi, Y.R. Access to pyridines via DMAP-catalyzed activation of α -chloro acetic ester to react with unsaturated imines. *Org. Chem. Front.* **2014**, *1*, 148–150. [\[CrossRef\]](#)
58. Young, C.M.; Taylor, J.E.; Smith, A.D. Evaluating aryl esters as bench-stable C (1)-ammonium enolate precursors in catalytic, enantioselective Michael addition–lactonisations. *Org. Biomol. Chem.* **2019**, *17*, 4747–4752. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Hartley, W.C.; O’Riordan, T.J.C.; Smith, A.D. Aryloxide-Promoted Catalyst Turnover in Lewis Base Organocatalysis. *Synthesis* **2017**, *49*, 3303–3310.
60. West, T.H.; Daniels, D.S.B.; Slawin, A.M.Z.; Smith, A.D. An isothiourea-catalyzed asymmetric [2, 3]-rearrangement of allylic ammonium ylides. *J. Am. Chem. Soc.* **2014**, *136*, 4476–4479. [\[CrossRef\]](#)
61. West, T.H.; Walden, D.M.; Taylor, J.E.; Brueckner, A.C.; Johnston, R.C.; Cheong, P.H.-Y.; Lloyd-Jones, G.C.; Smith, A.D. Catalytic Enantioselective [2, 3]-Rearrangements of Allylic Ammonium Ylides: A Mechanistic and Computational Study. *J. Am. Chem. Soc.* **2017**, *139*, 4366–4375. [\[CrossRef\]](#)
62. Spoehrle, S.S.M.; West, T.H.; Taylor, J.E.; Slawin, A.M.Z.; Smith, A.D. Tandem Palladium and Isothiourea Relay Catalysis: Enantioselective Synthesis of α -Amino Acid Derivatives via Allylic Amination and [2, 3]-Sigmatropic Rearrangement. *J. Am. Chem. Soc.* **2017**, *139*, 11895–11902. [\[CrossRef\]](#)
63. Kasten, K.; Slawin, A.M.Z.; Smith, A.D. Enantioselective Synthesis of β -Fluoro- β -aryl- α -aminopentenamides by Organocatalytic [2, 3]-Sigmatropic Rearrangement. *Org. Lett.* **2017**, *19*, 5182–5185. [\[CrossRef\]](#)
64. West, T.H.; Spoehrle, S.S.M.; Smith, A.D. Isothiourea-catalysed chemo- and enantioselective [2, 3]-sigmatropic rearrangements of N, N-diallyl allylic ammonium ylides. *Tetrahedron* **2017**, *73*, 4138–4149. [\[CrossRef\]](#)
65. Schwarz, K.J.; Amos, J.L.; Klein, J.C.; Do, D.T.; Snaddon, T.N. Uniting C1-ammonium enolates and transition metal electrophiles via cooperative catalysis: The direct asymmetric α -allylation of aryl acetic acid esters. *J. Am. Chem. Soc.* **2016**, *138*, 5214–5217. [\[CrossRef\]](#)
66. Schwarz, K.J.; Pearson, C.M.; Cintron-Rosado, G.A.; Liu, P.; Snaddon, T.N. Traversing Steric Limitations by Cooperative Lewis Base/Palladium Catalysis: An Enantioselective Synthesis of α -Branched Esters Using 2-Substituted Allyl Electrophiles. *Angew. Chem.* **2018**, *130*, 7926–7929. *Angew. Chem.* **2018**, *57*, 7800–7803. [\[CrossRef\]](#)

67. Scaggs, W.R.; Snaddon, T.N. Enantioselective α -Allylation of Acyclic Esters using B(pin)-Substituted Electrophiles: Independent Regulation of Stereocontrol Elements via Cooperative Pd/Lewis Base Catalysis. *Chem. Eur. J.* **2018**, *24*, 14378–14381. [\[CrossRef\]](#)
68. Fyfe, J.W.B.; Kabia, O.M.; Pearson, C.M.; Snaddon, T.N. Si-directed regiocontrol in asymmetric Pd-catalyzed allylic alkylations using C1-ammonium enolate nucleophiles. *Tetrahedron* **2018**, *74*, 5383–5391. [\[CrossRef\]](#)
69. Hutchings-Goetz, J.; Yang, C.; Snaddon, T.N. Enantioselective Syntheses of Strychnos and Chelidonium Alkaloids through Regio- and Stereocontrolled Cooperative Catalysis. *ACS Catal.* **2018**, *8*, 10537–10544. [\[CrossRef\]](#)
70. Pearson, C.M.; Fyfe, J.W.B.; Snaddon, T.N. A Regio- and Stereodivergent Synthesis of Homoallylic Amines by a One-Pot Cooperative-Catalysis-Based Allylic Alkylation/Hofmann Rearrangement Strategy. *Angew. Chem. Int. Ed.* **2019**, *58*, 10521–10527. *Angew. Chem.* **2019**, *131*, 10631–10637. [\[CrossRef\]](#)
71. Scaggs, W.R.; Scaggs, T.D.; Snaddon, T.N. An enantioselective synthesis of α -alkylated pyrroles via cooperative isothiourea/palladium catalysis. *Org. Biomol. Chem.* **2019**, *17*, 1787–1790. [\[CrossRef\]](#)
72. Schwarz, K.J.; Yang, C.; Fyfe, J.W.B.; Snaddon, T.N. Enantioselective α -Benzylation of Acyclic Esters Using π -Extended Electrophiles. *Angew. Chem. Int. Ed.* **2018**, *57*, 12102–12105. *Angew. Chem.* **2018**, *130*, 12278–12281. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Jiang, X.; Beiger, J.J.; Hartwig, J.F. Stereodivergent allylic substitutions with aryl acetic acid esters by synergistic iridium and Lewis base catalysis. *J. Am. Chem. Soc.* **2017**, *139*, 87–90. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Song, J.; Zhang, Z.J.; Gong, L.Z. Asymmetric [4+2] Annulation of C1 Ammonium Enolates with Copper-Allenylidenes. *Angew. Chem. Int. Ed.* **2017**, *56*, 5212–5216. *Angew. Chem.* **2017**, *129*, 5296–5300. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Song, J.; Zhang, Z.J.; Chen, S.S.; Fan, T.; Gong, L.Z. Lewis base/copper cooperatively catalyzed asymmetric α -amination of esters with diaziridinone. *J. Am. Chem. Soc.* **2018**, *140*, 3177–3180. [\[CrossRef\]](#) [\[PubMed\]](#)
76. Arokianathar, J.N.; Frost, A.B.; Slawin, A.M.Z.; Stead, D.; Smith, A.D. Isothiourea-Catalyzed Enantioselective Addition of 4-Nitrophenyl Esters to Iminium Ions. *ACS Catal.* **2018**, *8*, 1067–1075. [\[CrossRef\]](#)
77. Zhao, F.; Shu, C.; Young, C.M.; Carpenter-Warren, C.; Slawin, A.M.Z.; Smith, A.D. Enantioselective Synthesis of α -Aryl- β -2-Amino-Esters by Cooperative Isothiourea and Brønsted Acid Catalysis. *Angew. Chem. Int. Ed.* **2021**, *60*, 11892–11900. *Angew. Chem.* **2021**, *133*, 11999–12007. [\[CrossRef\]](#)
78. McLaughlin, C.; Slawin, A.M.Z.; Smith, A.D. Base-free Enantioselective C (1)-Ammonium Enolate Catalysis Exploiting Aryloxides: A Synthetic and Mechanistic Study. *Angew. Chem. Int. Ed.* **2019**, *131*, 15255–15263. *Angew. Chem.* **2019**, *58*, 15111–15119. [\[CrossRef\]](#)
79. McLaughlin, C.; Bitai, J.; Barber, L.; Slawin, A.M.Z.; Smith, A.D. Catalytic enantioselective synthesis of 1, 4-dihydropyridines via the addition of C (1)-ammonium enolates to pyridinium salts. *Chem. Sci.* **2021**. accepted for publication. [\[CrossRef\]](#)
80. Stockhammer, L.; Weinzierl, D.; Bögl, T.; Waser, M. Enantioselective α -Chlorination Reactions of in Situ Generated C1 Ammonium Enolates under Base-Free Conditions. *Org. Lett.* **2021**, *23*, 6143–6147. [\[CrossRef\]](#)
81. Kim, B.; Kim, Y.; Lee, S.Y. Stereodivergent Carbon–Carbon Bond Formation between Iminium and Enolate Intermediates by Synergistic Organocatalysis. *J. Am. Chem. Soc.* **2021**, *143*, 73–79.
82. Yuan, S.; Liao, C.; Zheng, W.-H. Paracyclophane-Based Isothiourea-Catalyzed Highly Enantioselective α -Fluorination of Carboxylic Acids. *Org. Lett.* **2021**, *23*, 4142–4146. [\[CrossRef\]](#)
83. Crystallographic data for (2S,1'R)-19 (CCDC 1992504), Cambridge Crystallographic Data Centre.
84. The relative and absolute configuration of all other products was assigned by analogy, with consistent diagnostic ^1H NMR signals of the proton at C(2) and C(1') stereocentres, indicating the same major diastereoisomer in each case.
85. Morrill, L.C.; Douglas, J.; Lebl, T.; Slawin, A.M.Z.; Fox, D.J.; Smith, A.D. Isothiourea-mediated asymmetric Michael-lactonisation of trifluoromethylenones: A synthetic and mechanistic study. *Chem. Sci.* **2013**, *4*, 4146–4155. [\[CrossRef\]](#)
86. Matviitsuk, A.; Greenhalgh, M.D.; Antúnez, D.-J.B.; Slawin, A.M.Z.; Smith, A.D. Aryloxide-Facilitated Catalyst Turnover in Enantioselective α , β -Unsaturated Acyl Ammonium Catalysis. *Angew. Chem. Int. Ed.* **2017**, *56*, 12282–12287. *Angew. Chem.* **2017**, *129*, 12450–12455. [\[CrossRef\]](#)
87. Catalyst turnover could also be envisaged through intramolecular nucleophilic displacement by the pendant phenol to give a cyclobutanone intermediate via a formal [2+2]-cycloaddition, which may then be ring-opened by *para*-nitrophenoxide to give the final product. Based on the high steric congestion of this proposed cyclobutanone intermediate, the mechanism given in Scheme 5 is considered more probable.
88. Liu, P.; Yang, X.; Birman, V.B.; Houk, K.N. Origin of enantioselectivity in benzotetramisole-catalyzed dynamic kinetic resolution of azlactones. *Org. Lett.* **2012**, *14*, 3288–3291. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Abbasov, M.E.; Hudson, B.M.; Tantillo, D.J.; Romo, D. Acylammonium salts as dienophiles in Diels–Alder/lactonization organocascades. *J. Am. Chem. Soc.* **2014**, *136*, 4492–4495. [\[CrossRef\]](#) [\[PubMed\]](#)
90. Robinson, E.R.T.; Walden, D.M.; Fallan, C.; Greenhalgh, M.D.; Cheong, P.H.-Y.; Smith, A.D. Non-bonding 1,5-S \cdots O interactions govern chemo- and enantioselectivity in isothiourea-catalyzed annulations of benzazoles. *Chem. Sci.* **2016**, *7*, 6919–6927. [\[CrossRef\]](#) [\[PubMed\]](#)
91. Greenhalgh, M.D.; Smith, S.M.; Walden, D.M.; Taylor, J.E.; Brice, Z.; Robinson, E.R.T.; Fallan, C.; Cordes, D.B.; Slawin, A.M.Z.; Richardson, H.C.; et al. AC=O \cdots Isothiuronium Interaction Dictates Enantiodiscrimination in Acylative Kinetic Resolutions of Tertiary Heterocyclic Alcohols. *Angew. Chem. Int. Ed.* **2018**, *57*, 3200–3206. *Angew. Chem.* **2018**, *130*, 3254–3260. [\[CrossRef\]](#)
92. Young, C.M.; Elmi, A.; Pascoe, D.J.; Morris, R.K.; McLaughlin, C.; Woods, A.M.; Frost, A.B.; de la Houpliere, A.; Ling, K.B.; Smith, T.K.; et al. The Importance of 1, 5-Oxygen Chalcogen Interactions in Enantioselective Isochalcogenourea Catalysis. *Angew. Chem.* **2020**, *132*, 3734–3739. *Angew. Chem.* **2020**, *59*, 3705–3710. [\[CrossRef\]](#)

-
93. Pascoe, D.J.; Ling, K.B.; Cockroft, S.L. The origin of chalcogen-bonding interactions. *J. Am. Chem. Soc.* **2017**, *139*, 15160–15167. [[CrossRef](#)]
 94. Nagao, Y.; Miyamoto, S.; Miyamoto, M.; Takeshige, H.; Hayashi, K.; Sano, S.; Shiro, M.; Yamaguchi, K.; Sei, Y. Highly Stereoselective Asymmetric Pummerer Reactions That Incorporate Intermolecular and Intramolecular Nonbonded S···O Interactions. *J. Am. Chem. Soc.* **2006**, *128*, 9722–9729. [[CrossRef](#)]
 95. Beno, B.R.; Yeung, K.-S.; Bartberger, M.D.; Pennington, L.D.; Meanwell, N.A. A survey of the role of noncovalent sulfur interactions in drug design. *J. Med. Chem.* **2015**, *58*, 4383–4438. [[CrossRef](#)]
 96. Breugst, M.; Koenig, J.J. σ -Hole Interactions in Catalysis. *Eur. J. Org. Chem.* **2020**, *34*, 5473–5487. [[CrossRef](#)]
 97. Bleiholder, C.; Gleiter, R.; Werz, D.B.; Köppel, H. Theoretical Investigations on Heteronuclear Chalcogen–Chalcogen Interactions: On the Nature of Weak Bonds between Chalcogen Centers. *Inorg. Chem.* **2007**, *46*, 2249–2260. [[CrossRef](#)]
 98. Gleiter, R.; Haberhauer, G.; Werz, D.B.; Rominger, F. From noncovalent chalcogen–chalcogen interactions to supramolecular aggregates: Experiments and calculations. *Bleiholder. Chem. Rev.* **2018**, *118*, 2010–2041. [[CrossRef](#)] [[PubMed](#)]
 99. Benz, S.; López-Andarias, J.; Mareda, J.; Sakai, N.; Matile, S. Catalysis with chalcogen bonds. *Angew. Chem.* **2017**, *129*, 830–833. *Angew. Chem.* **2017**, *56*, 812–815. [[CrossRef](#)]
 100. Wonner, P.; Vogel, L.; Düser, M.; Gomes, L.; Kniep, F.; Mallick, B.; Werz, D.B.; Huber, S.M. Carbon–Halogen Bond Activation by Selenium-Based Chalcogen Bonding. *Angew. Chem. Int. Ed.* **2017**, *56*, 12009–12012. *Angew. Chem.* **2017**, *129*, 12172–12176. [[CrossRef](#)] [[PubMed](#)]
 101. Wonner, P.; Vogel, L.; Kniep, F.; Huber, S.M. Catalytic Carbon–Chlorine Bond Activation by Selenium-Based Chalcogen Bond Donors. *Chem. Eur. J.* **2017**, *23*, 16972–16975. [[CrossRef](#)]
 102. Wonner, P.; Dreger, A.; Vogel, L.; Engelage, E.; Huber, S.M. Chalcogen Bonding Catalysis of a Nitro-Michael Reaction. *Angew. Chem. Int. Ed.* **2019**, *58*, 16923–16927. *Angew. Chem.* **2019**, *131*, 17079–17083. [[CrossRef](#)]
 103. Wang, W.; Zhu, H.; Liu, S.; Zhao, Z.; Zhang, L.; Hao, J.; Wang, Y. Chalcogen–chalcogen bonding catalysis enables assembly of discrete molecules. *J. Am. Chem. Soc.* **2019**, *141*, 9175–9179. [[CrossRef](#)]
 104. Wang, W.; Zhu, H.; Feng, L.; Yu, Q.; Hao, J.; Zhu, R.; Wang, Y. Dual Chalcogen–Chalcogen Bonding Catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 3117–3124. [[CrossRef](#)]